

Povidone lodine

10 % Vaginal Douche

USED FOR

- Vaginitis infections
- Infection preventer
- Personal Hygiene

500 ml



Drug Facts:

Active ingredient:

Povidone Iodine 10 %, PEG, Poloxamer, Citric-acid, Phosphate buffer

Uses:

- For personal Hygiene
- For treatment of vaginal infection including nonspecific vaginitis and vulvovaginitis associated with trichomonas vaginalis

Warnings: For external use only

Do not use

o The product is spermicidal and should not be used when consumption is desired

Stop use and ask a doctor:

- · If condition persists or gets worse
- If redness develop and persist for over 72 hours

Keep our of reach of children

If swallowed get medical help or contact a poison control center right away

Direction of use:

- To be used together with a douching kit
- Routine cleaning and deodorizing: Dilute 6 to 12 ml of Povidone solution in 1 Liter of warm water no more than once or twice a week.
- Therapeutic use: Dilute 25 ml of Povidone solution in 1 liter of warm water, treatment should be continued for 14 days and should continue during menstruation.
- Since Povidone may stain synthetic fabric, use of a feminine hygiene pad is recommended.

Produced by:- Amshaj manufacturing plc, Addis Ababa, Ethiopia www.jourilabs.com +251-11-5580481/2

